



Review Article

## Silicosis- An Incurable Lung Disease

Shukla Prabodh<sup>1\*</sup>, Shukla Padmini<sup>1</sup>, Gopalakrishna B.<sup>2</sup>

1 Pranveer Singh Institute of Technology, Kanpur, (U.P.), India.

2 R. R. College of Pharmacy, Chikbanawara, Bangalore, India.

### ABSTRACT

Silica is the second most common element on the surface of the earth. It is found in sand, many rocks such as granite, sandstone, flint and slate, and in some coal and metallic ores. The cutting, breaking, crushing, drilling, grinding, or abrasive blasting of these materials may produce fine silica dust. Silicosis is a form of occupational lung disease caused by inhalation of crystalline silica dust, and is marked by inflammation and scarring in forms of nodular lesions in the upper lobes of the lungs. Silicosis is due to deposition of fine dust (less than 1 $\mu$ m in diameter) containing crystalline silicon dioxide in the form of alpha-quartz, cristobalite, or tridymite. Silicosis is characterized by shortness of breath, fever, and cyanosis. It may often be misdiagnosed as pulmonary edema (fluid in the lungs), pneumonia, or tuberculosis. Treatment options currently focus on alleviating the symptoms and preventing complication. So here an attempt has been made to attract the attention of doctors, researchers, NGO's to develop a proper medication and vaccine kit for the complete treatment of this disease.

**Key words:** Silicosis, silica dust, treatment

**Corresponding Author:** Shukla Prabodh, EWS-915, Avas Vikas Colony-3, Kalyanpur Panki Road, Kanpur (U.P.) - 208017.

E-mail: shuklapp2000@gmail.com

**Article Info: Date received: 17 Jan. 2010**

**Date accepted : 21 March 2010**

### INTRODUCTION

Silicosis, known as Grinder's disease and Potter's rot, is a form of occupational lung disease caused by inhalation of crystalline silica dust, and is marked by inflammation and scarring in forms of nodular lesions in the upper lobes of the lungs[1,2]. Silica is the second most common element on the surface of the earth. It is found in sand, many rocks such as granite, sandstone, flint and slate, and in some coal and metallic ores. The cutting,

breaking, crushing, drilling, grinding, or abrasive blasting of these materials may produce fine silica dust. It can also be in soil, mortar, plaster, and shingles. Silicosis is due to deposition of fine dust (less than 1 micrometer in diameter) containing crystalline silicon dioxide in the form of alpha-quartz, cristobalite, or tridymite. Silicosis is the most common occupational lung disease worldwide; it occurs everywhere but is

especially common in developing countries [3, 4].

## **EPIDEMIOLOGY**

In India, a prevalence of 55% was found in one group of workers, many of them very young, engaged in the quarrying of shale sedimentary rock and subsequent work in small, poorly ventilated sheds. Studies on silicotic pencil workers in Central India demonstrated high mortality rates; the mean age at death was 35 years and the mean duration of the exposure was 12 years [5,6]. During the period 1991 to 1995, China recorded more than 5, 00 000 cases of silicosis, with around 6,000 new cases and more than 24,000 deaths occurring each year mostly among older workers [7].

### **Types of silicosis**

Classification of silicosis is made according to the disease's severity, onset, and rapidity of progression [8, 9,10]. These include: Chronic silicosis occurs after 15–20 years of exposure to moderate to low levels of silica dust. Chronic silicosis itself is further subdivided into simple and complicated silicoses. This is the most common type of silicosis. Patients with this type of silicosis may not have obvious symptoms, so a chest X-ray is necessary to determine if there is lung damage.

### **Asymptomatic silicosis**

Early cases of the disease do not present any symptoms.

### **Accelerated silicosis**

Silicosis develops in 5–10 years after high exposure to silica dust. Symptoms include severe shortness of breath, weakness, and weight loss.

### **Acute silicosis**

Silicosis that develops in few months to 2 years after exposure to very high concentrations of silica dust. Symptoms of acute silicosis include severe disabling shortness of breath, weakness, and weight loss, often leading to death.

## **SYMPTOMS**

Silicosis (especially the acute form) is characterized by shortness of breath, fever, and cyanosis (bluish skin). It may often be misdiagnosed as pulmonary edema (fluid in the lungs), pneumonia, or tuberculosis [11,12,13].

- Dyspnea exacerbated by exertion
- Dry or severe cough, often persistent and accompanied by hoarseness of the throat
- Fatigue
- Tachypnea (rapid breathing) which is often labored
- Loss of appetite
- Chest pain
- Fever

Gradual dark shallow in nail rifts eventually leading to cracks as protein fibers within nail beds are destroyed. In advanced cases, the following may also occur:

- Cyanosis
- Cor pulmonale
- Respiratory insufficiency

Silicotuberculosis- Patients with silicosis are particularly Susceptible to tuberculosis infection known as silico tuberculosis. The

reason for the increased risk 10–30 fold increased incidence is not well understood. It is thought that silica damages pulmonary macrophages, inhibiting their ability to kill mycobacteria.

### **PATHOPHYSIOLOGY**

When small silica dust particles are inhaled, they can embed themselves deeply into the tiny alveolar sacs and ducts in the lungs, where oxygen and carbon dioxide gases are exchanged. There, the lungs cannot clear out the dust by mucous or coughing [13]. When fine particles of silica dust are deposited in the lungs, macrophages that ingest the dust particles will set off an inflammation response by releasing tumor necrosis factors, interleukin-1, leukotriene B4 and other cytokines. In turn, these stimulate fibroblasts to proliferate and produce collagen around the silica particle, thus resulting in fibrosis and the formation of the nodular lesions. The inflammatory effects of crystalline silica are apparently mediated by the Nalp3 inflammasome[14]. Furthermore, the surface of silicon dust can generate silicon-based radicals that lead to the production of hydroxyl and oxygen radicals, as well as hydrogen peroxide, which can inflict damage to the surrounding cells[15].

### **DIAGNOSIS**

Characteristic lung tissue pathology in nodular silicosis consists of fibrotic nodules with concentric "onion-skinned" arrangement of collagen fibers, central hyalinization, and a cellular peripheral zone, with lightly birefringent particles seen under polarized light.[16] In acute silicosis, microscopic pathology shows a periodic acid-

Schiff positive alveolar exudate (alveolar lipoproteinosis) and a cellular infiltrate of the alveolar walls. Physical check up will reveal decreased chest expansion and abnormal breath sounds. Pulmonary function test will reveal reduced lung capacity. [17,18] A computed tomography or CT scan can also provide a more detailed analyses of the nodules, and can reveal cavitation due to concomitant mycobacterial infection. Silicosis is an irreversible condition with no cure. Chest x-ray will confirm the presence of nodules in the lungs, especially in the upper lobes. Typically, it will also reveal eggshell calcification of the hilar lymph nodes. In rare cases, pulmonary nodules may also be calcified. In advanced cases of silicosis, coalescence of nodules may show up as large masses.[19,20]

### **TREATMENT**

Treatment options currently focus on alleviating the symptoms and preventing complications. [21,22] These include:

1. Stopping further exposure to silica and other lung irritants, including tobacco smoking.
2. Cough suppressants.
3. Antibiotics and antitubercular agents to prevent tuberculosis. These include isoniazid, rifampin, and pyrazinamide.
4. Chest physiotherapy to help the bronchial drainage of mucus.
5. Oxygen administration to avoid hypoxemia.
6. Bronchodilators to facilitate breathing.
7. Lung transplantation to replace the damaged lung tissue is the most effective treatment, but is associated with severe risks of its own.

9. Whole-lung lavage
10. Inhalation of powdered aluminium, d-penicillamine and polyvinyl pyridine-N-oxide.
11. Corticosteroid therapy.
12. The herbal extract tetrandine may slow progression of silicosis.[23,24]

### **PREVENTION**

The following precaution should be taken by the workers to prevent from silicosis[25,26]. Wear dust masks Do not smoke Respirator should be provided from the organizations Water spray is often used where dust emanates Dust can also be controlled through dry air filtering Identify work-place activities that produce crystalline silica dust and then to eliminate or control the dust.



Fig. 1 X-Ray report of silicotic patient

### **CONCLUSION**

From the above study it can be concluded that the silicosis is an incurable disease for both coal and stone mine workers, which is caused by inhalation of silica dust particles. By this study an attempt has been made to attract the attention of doctors, researchers, NGO's to develop a proper medication and vaccine kit for the complete treatment of this

disease. It is also suggested that the precautions should be taken to get rid from this disease because precaution is better than cure.

### **REFERENCES**

1. Mulloy K.B., Silica exposure and systemic vasculitis. *Environ Health Perspect.* 2003, 111(16), 1933-8.
2. Banks D.E., Balaan M., Wang M.L., Silicosis in the 1990s, revisited. *Chest.* 1997, 111(4), 837-8.
3. Rosenman K.D., Moore-Fuller M., Reilly M.J., Connective tissue disease and silicosis. *Am J. Ind. Med.* 1999, 35(4), 375-81.
4. Caplan A., Payne R.B., Withley J.L., A broadened concept of Caplan's syndrome related to rheumatoid factors. *Thorax.* 1962, 17, 205-209.
5. Graham W.G.B., Quartz and silicosis. In: Banks D, Parker J, eds. *Occupational Lung Disease: An International Perspective.* New York, NY: Chapman & Hall Medical; 1998, 191-212.
6. Hertzberg V.S., Rosenman K.D., Reilly M.J., Rice C.H., Effect of occupational silica exposure on pulmonary function. *Chest.* 2002, 122(2),721-8.
7. Wagner G.R., Screening and surveillance of workers exposed to mineral dusts. Geneva, World Health Organization. 1996.
8. Gamble J.F., Hessel P.A., Nicolich M., Relationship between silicosis and lung

- function., Scand J Work Environ Health. 2004, 30(1), 5-20.
9. [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&list\\_uids=8561170&dopt=Citation](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&list_uids=8561170&dopt=Citation).
  10. Ghio A.J., Kennedy T.P., Schapira R.M., Crumbliss A.L., Hoidal J.R., Hypothesis: silicosis is lung disease after silicate inhalation caused by oxidant generation., Lancet. 1990, 336(8721), 967-9.
  11. Attfield M.D., Costello J., Quantitative exposure-response for silica dust and lung cancer in vermont granite workers. Am., J. Ind. Med. Feb 2004, 45(2), 129-38.
  12. Centers for Disease Control and Prevention. Silicosis in dental laboratory technicians--five states, 1994-2000. MMWR Morb Mortal Wkly Rep. 2004, 53(9), 195-7.
  13. Arakawa H., Honma K., Saito Y., Shida H., Morikubo H., Suganuma N., Pleural disease in silicosis: pleural thickening, effusion, and invagination. Radiology. 2005, 236(2), 685-93.
  14. Cassel S.L., Eisenbarth S.C., Iyer S.S., et al., The Nalp3 inflammasome is essential for the development of silicosis. Proc. Natl. Acad. Sci., 2008, 105, 9035.
  15. Corbett E.L., Churchyard G.J., Clayton T., Herselman P., Williams B., Hayes R., Risk factors for pulmonary mycobacterial disease in South African gold miners. A case-control study. Am J Respir Crit Care Med. 1999, 159(1), 94-9.
  16. International Journal of Pharmacy Research 2010, 1 (1), 14-19
  17. American Thoracic Society, Targeted tuberculin testing and treatment of latent tuberculosis infection, Am. J. Respir. Crit. Care. Med. 2000, 161(4 Pt 2), S221-47.
  18. Corbett E.L., Murray J., Churchyard G.J., Herselman P.C., Clayton T.C., De Cock K.M., Use of miniradiographs to detect silicosis. Comparison of radiological with autopsy findings. Am J Respir Crit Care Med. 1999, 160(6), 2012-7.
  19. <http://www.osha.gov/SLTC/silicacrystalline/index.html>.
  20. <http://www.who.int/mediacentre/factsheets/fs238/en/>
  21. Lapp N.L., Pulmonary function and disability. In: Banks D, Parker J, eds. Occupational Lung Disease: An International Perspective. 1998. New York, NY: Chapman & Hall Medical; 1998,17-34.
  22. Goodman G.B., Kaplan P.D., Stachura I., Castranova V., Pales W.H., Lapp N.L., Acute silicosis responding to corticosteroid therapy. Chest. 1992, 101(2), 366-70.
  23. Sharma S.K., Pande J.N., Verma K., Effect of prednisolone treatment in chronic silicosis. Am Rev Respir Dis. 1991, 143(4 Pt 1), 814-21.
  24. McDonald J.C., McDonald A.D., Hughes J.M., Rando R.J., Weill H., Mortality from lung and kidney disease in a cohort of

- North American industrial sand workers: an update. *Ann Occup Hyg.* 2005, 49(5), 367-73.
25. National Archives and Records Administration. United States code of federal regulations. 1910.1000. Office of the Federal Register, National Archives and Records Administration;1994.
26. Rimal B., Greenberg A.K., Rom W.N., Basic pathogenetic mechanisms in silicosis: current understanding. *Curr. Opin. Pulm. Med.* 2005, 11(2),169-73.
27. Valiante D.J., Schill D.P., Rosenman K.D., Socie E. Highway repair: a new silicosis threat. *Am. J. Public. Health.* 2004, 94(5), 876-80.